

CERN-TH.6464/92

April, 1992

A PROPOSAL FOR $D > 1$ STRINGS?

L. ALVAREZ-GAUMÉ

J.L.F. BARBÓN[†]*Theory Division, CERN**CH-1211 Geneva 23, Switzerland*

ABSTRACT

Using the reduced formulation of large- N Quantum Field Theories we study strings in space-time dimensions higher than one. Some preliminary results concerning the possible string susceptibilities and general properties of the model are presented.

★ Talk presented at the Trieste Spring Workshop. April 8-11, 1992.

† Permanent address: Departamento de Física Teórica, Universidad Autónoma de Madrid, Canto Blanco, Madrid. Spain.

1. INTRODUCTION

In this talk I would like to report on some preliminary work done in collaboration with J.L.F. Barbón on the application of large- N reduced models [1,2,3] to the study of non-critical strings in dimensions higher than one. This is a rather challenging problem, and at the moment we can only make few general remarks on a proposal to construct these theories. The first lattice formulations of string theories in arbitrary dimensions appeared in the mid-eighties [4,5,6,7], and there was a good deal of activity in the solution of some two-dimensional models in random triangulated surfaces [10]. A breakthrough took place with the work in ref. [8] which provided a description in the continuum of the coupling of minimal conformal models to two-dimensional gravity in the light-cone gauge. The subsequent formulation of these models in the conformal gauge and the generalization of the results to surfaces of arbitrary topology appeared in [9]. With the discovery of the double scaling limit [11] we began to understand some of the perturbative and non-perturbative properties of non-critical strings in dimensions below or equal to one. Little progress has been made however in going beyond $d = 1$ apart from some evidence that it is possible to construct Liouville Quantum Field Theory in some especial dimensions $d = 7, 13, 19$.

Some interesting problems involving the presence of external have also been studied [13,14,15,16]. In particular the work in [16] attempted to generalize the standard one-matrix model to include the presence of local curvatures in the triangulated surfaces described by the large- N limit. We will make contact with this work later in this lecture.

The outline of this talk is as follows: In section one we present the problem of strings in arbitrary dimensions in general. In section two we briefly review the definition of the reduced models in the large- N description of matrix field theories, and the general features of the effective one-matrix model formulation of strings propagating in D -dimensional flat space-time. In section three we present the general features of the model and study the simplest possible approximation

leading to a polymer phase. In section four we study more precise approximations as well as the critical properties of some effective actions suggested from our model. We show that for these effective actions we can obtain positive string susceptibilities $\gamma_{\text{st}} = n/(n + m + 1)$ with n, m arbitrary positive integers. We compute correlators of macroscopic loop operators on the surface and show that only for a subclass of critical points ($n = 1$) is it possible to define macroscopic loops in the continuum limit. Furthermore the continuum limit of these theories contains an extra state with respect to the standard pure gravity case, representing the breaking of the surface in two pieces. This extra state is probably the shadow of the tachyon in the approximations to our model that we study. We also explore some of the scaling operators appearing in truncations of the reduced model. This section contains our main results. The outlook and possible future avenues to be investigated, together with the conclusions appear in section five.

2. LARGE- N REDUCED THEORIES AND MATRIX MODELS

In the study of strings propagating in flat D -dimensional space-time, we represent the sum over two-dimensional metrics in Polyakov's approach to string theory [17] by a sum over tringulated surfaces. The quantity we would like to evaluate [4,6,7] is

$$Z = \sum_g \kappa^{2(g-1)} \sum_T \frac{e^{-\mu|T|}}{|n(T)|} \int \prod_{i \in T_0} \sigma_i^\alpha d^D X_i \prod_{\langle ij \rangle \in T_1} G(X_i - X_j) \quad (2.1)$$

Where g is the genus of the triangulation. For a given triangulation T , T_0, T_1, T_2 are respectively the sets of vertices, edges and faces of T , $n(T)$ is the order of the symmetry group of T , $|T|$ is the total area of T counting that every triangle has unit area. μ is the bare two-dimensional cosmological constant, X_i describes the embedding of the triangulation into D -dimensional flat space-time and $G(X - Y)$ is the propagator factor for each link. We have included also the local volume factor σ_i^α at site i to represent the effect of local curvature. Usually we take $\alpha = D/2$,

but it is more convenient to leave this exponent arbitrary to include the effect of local curvature terms on the world-sheet. For a triangulation, if q_i is the local coordination number at site i (the number of triangles sharing this vertex),

$$\sigma_i = \frac{q_i}{3},$$

and the volume factor in the measure can be shown to be related in the continuum to an expansion of the form:

$$\sum_i \ln \sigma_i = c_0 + c_1 \int \sqrt{g} R + c_2 \int \sqrt{g} R^2 \dots,$$

thus representing some of the effects of world-sheet curvature. The general features of the phase diagram in the (D, α) plane were studied for example in [18]. In the standard Polyakov formulation, the propagator is Gaussian,

$$G(X) = e^{-X^2/2}$$

By standard large-N analysis the sum (2.1) can be transformed [6,7] into the large- N expansion of a matrix field theory in D -dimensions

$$Z = \lim_{N \rightarrow \infty} \log \int D\phi \exp \left(-N \int d^D X \text{Tr} \left(\frac{1}{2} \phi G^{-1} \phi + \frac{1}{3} g \phi^3 \right) \right) \quad (2.2)$$

It is difficult to proceed very far with Gaussian propagators. If we work in dimensions $D < 6$ it should not matter whether we replace the Gaussian propagator by the Feynman propagator. There is numerical evidence that this change does not affect the critical properties of the theory below six dimensions [19]. Notice that in this construction the exponent α in (2.1) is set to zero. If we want to include the effect of local world-sheet curvature, we can follow [16] and change the kinetic term in (2.2). The two-dimensional cosmological constant is here represented by

g , $g = e^{-\mu}$. To summarize, we want to study the critical properties of the action:

$$Z = \int D\phi(X) \exp \left(-N \int d^D X \text{Tr} \left(\frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \frac{g}{3} \phi^3 \right) \right) , \quad (2.3)$$

or

$$Z = \int D\phi(X) \exp \left(-N \int d^D X \text{Tr} \left(\frac{1}{2} A \partial_\mu \phi A \partial^\mu \phi + \frac{g}{3} \phi^3 \right) \right) , \quad (2.4)$$

where A is a constant $N \times N$ matrix and ϕ is an $N \times N$ matrix field. The reason why the A -matrix in the kinetic term simulates the effect of local curvature can be seen by writing the propagator in (2.4) explicitly

$$\langle \phi_{ij}(X) \phi_{kl}(Y) \rangle = A_{jk}^{-1} A_{li}^{-1} G(X - Y).$$

If we ignore for the time being the propagator factor G , for every closed index loop in a generic ϕ^3 graph made of q propagators, we obtain a contribution of $\text{tr} A^{-q}$. Since the ϕ^3 graphs are dual to triangulation, this means that we are associating a curvature factor of $\text{tr} A^{-q}$ to the vertex dual to the face considered. In this way we can simulate the presence of the σ^α term in the measure in (2.2). We are interested in particular in the computation of the string susceptibility exponent

$$\chi \sim \frac{\partial^2 F}{\partial g^2} \sim (g - g_c)^{-\gamma_{\text{st}}}, \quad (2.5)$$

where F is the free energy of the system, and g_c is the critical value of g indicating the location of the critical point. Apart from the string susceptibility we would also like to compute the spectrum of scaling operators and the properties of the quantum geometry implied by (2.3)(2.4).

To simplify the arguments we concentrate on the study of planar configurations (spherical topologies). From work dating back to the late seventies [20], and due to the factorization properties of the leading large- N approximation, the planar limit is dominated by a single constant configuration; Witten's master field (a

master orbit in the case of gauge theories). This idea is explicitly realized in 0-dimensional matrix models [21] and in lattice gauge theories one can think of the reduced Eguchi-Kawai (EK)[1] or Twisted Eguchi-Kawai (TEK)[2] models as explicit descriptions of the master field. We now briefly describe the main ideas behind the reduced models. Further details can be found in the previous references and in the review article [3]. These models were originally formulated to describe the large- N properties of lattice gauge theories.

Due to the factorization property in the large- N limit, the Schwinger-Dyson equations for Wilson-loops form a closed system of equations in the planar limit. For gauge theories we get an infinite set of polynomial equations for the Wilson-loops [22]. In terms of the standard link variables $U_\mu(x)$, the EK proposal consists of making them independent of x , $U_\mu(x) \mapsto U_\mu$, and the action of the reduced model becomes

$$S_{EK} = \frac{1}{\text{Vol.}} S(U_\mu(x) \mapsto U_\mu). \quad (2.6)$$

Some of the properties of this action are:

1). We obtain a theory on a single hypercube with periodic boundary conditions. This is an important simplification of the problem.

2). In the case of gauge theories, the gauge symmetry becomes a global symmetry, $U_\mu \mapsto \Omega U_\mu \Omega^{-1}$.

3). There is an extra $U(1)^D$ symmetry in the case of $U(N)$ lattice gauge theory $U_\mu \mapsto e^{i\theta_\mu} U_\mu$. In the case of $SU(N)$ the symmetry becomes Z_N^D .

4). It is possible to show that the planar Schwinger-Dyson equations following from the reduced action coincide with those obtained from the original Wilson theory as long as open loop expectation values vanish. This can be shown to be true at strong coupling, but it does not hold at weak coupling [23]. Without this problem we would have a beautiful implementation of the Master Field idea, because the loop equations for the 1-site EK model are identical with the standard ones. The problem is connected with the fact that at weak coupling the extra

$U(1)^D$ symmetry is broken. Open loops have non-trivial charge with respect to this symmetry. Only if the symmetry remains unbroken are we guaranteed to maintain the loop equations without extraneous terms.

The resolution of this problem motivated the formulation of the TEK model [2]. It is inspired on 'tHooft's use of twisted boundary conditions in gauge theories [24]. For any matrix theory the TEK prescription is exceedingly simple. We reduce according to

$$\phi(x) \mapsto D(x)\phi D(x)^{-1}, \quad (2.7)$$

where $D(x)$ is a projective representation of the D -dimensional lattice translation group. We have to choose a set of D N -dimensional matrices Γ_μ such that:

$$D(x) = \prod_{\mu} \Gamma_{\mu}^{x_{\mu}} \quad x = (x_1, x_2, \dots, x_D) \quad x_i \in Z, \quad (2.8)$$

since $D(x)$ has an adjoint action on the fields ϕ , the Γ_μ 's are required to commute only up to an element of the center of $SU(N)$,

$$\Gamma_\mu \Gamma_\nu = Z_{\nu\mu} \Gamma_\nu \Gamma_\mu$$

$$Z_{\mu\nu} = e^{2\pi i n_{\mu\nu}/N}, \quad (2.9)$$

and the integers $n_{\mu\nu}$ are defined mod N . The reduced action prescription is now

$$S_{TEK} = \frac{1}{\text{Vol.}} S(\phi(x) \mapsto D(x)\phi D(x)^{-1}), \quad (2.10)$$

and similarly for expectation values.

To avoid open loop expectation values and a mismatch between the original and reduced Schwinger-Dyson equations, the matrices Γ_μ must verify some conditions. In particular they should generate an irreducible representation of the group of lattice translations. For a lattice with L^D sites this requires $N = L^{D/2}$. Hence

$N = L$ for $D = 2$ and $N = L^2$ for $D = 4$. Hence, if we want to simulate a two-dimensional lattice with 64×64 sites it suffices to consider the group $SU(64)$. The choice of twist matrix $n_{\mu\nu}$ depends on the dimensionality. In two-dimensions, the simplest choice is given by

$$n_{\mu\nu} = \epsilon_{\mu\nu} \quad (2.11)$$

The explicit form of the four and higher dimensional twists can be found in the quoted literature.

We now set aside lattice gauge theory and return to our problem. The twisted reduced version of (2.3) becomes:

$$S = \frac{1}{2} \sum_{\mu} Tr(\Gamma_{\mu} \phi \Gamma_{\mu}^{-1} - \phi)^2 + Tr V(\phi). \quad (2.12)$$

The equivalence of the planar approximation to (2.3) with (2.12) follows from earlier work on reduced models [2,25]. For simplicity we consider the two-dimensional case. First, one can always write a hermitean matrix in the Γ_{μ} basis

$$A(q) = \Gamma_1^{k_1} \Gamma_2^{k_2} \quad k_{\mu} = \epsilon_{\mu\nu} q_{\nu},$$

with q_{ν} defined modulo N . We can expand the hermitean matrix ϕ in the Γ -basis according to

$$\phi = \sum \phi_q A(q).$$

Some useful properties of the A -matrices are:

$$A(q)^{\dagger} = A(-q) e^{\frac{2\pi i}{N} \langle k|k \rangle} \quad \langle k^i | k^j \rangle = \sum_{\mu < \nu} n_{\mu\nu} k_{\nu}^i k_{\mu}^j,$$

and

$$\Gamma_{\mu} A(q) \Gamma_{\mu}^{\dagger} = e^{2\pi i q_{\mu} / L} A(q),$$

explicitly showing how the Γ 's implement the lattice translation group. Further-

more

$$Tr A(q_1)^\dagger A(q_2) = N \delta_L(q_1 - q_2),$$

where the δ -function is defined mod L . Using these properties, the kinetic term of our action becomes,

$$\begin{aligned} & \frac{1}{2} \sum_{\mu} Tr(\Gamma_{\mu} \phi \Gamma_{\mu}^{-1} - \phi)^2 + \frac{1}{2} m^2 Tr \phi^2 \sim \\ & \sum_q \left[\frac{1}{2} m^2 + \sum_{\mu} (1 - \cos(\frac{2\pi q_{\mu}}{L})) \right] \phi_q \phi_{-q} e^{-2\pi i \langle k|k \rangle / N}, \end{aligned} \quad (2.13)$$

coinciding with the standard lattice propagator of a scalar field on a size L D -dimensional lattice up to the last phase factor in (2.13). The role of this and similar phase factors appearing in the vertices of the theory is to restore the topological expansion in large- N . We have now obtained an effective field theory with scalar variables ϕ_q . The phase factors keep track of the fact that the theory came from a matrix field theory and that the expansion in powers of $1/N$ can be represented in terms of a topological expansion where the leading order corresponds to spherical topologies, and higher orders come from surfaces of increasing Euler number. The phases cancel completely only for planar graphs, and for higher topologies the lack of cancellation leads to suppressions by powers of $1/N^2$. Note that in (2.13) one shows that the space-time degrees of freedom of the original theory are coded in the Fourier transforms of the “internal” $SU(N)$ indices.

The construction we just carried out works only for even dimensions. For odd-dimensions we can leave one of the dimensions unreduced, and apply the reduction prescription to the remaining even number of dimensions. This yields an action

$$\begin{aligned} S = & \int dt \frac{1}{2} Tr \dot{\phi}(t)^2 + \frac{1}{2} \int dt Tr(\Gamma_{\mu} \phi(t) \Gamma_{\mu}^{-1} - \phi)^2 \\ & + \int dt Tr V(\phi(t)). \end{aligned}$$

We can consider a slightly more general model by including a general coupling in

the hopping term:

$$\begin{aligned}
-S(\phi) &= \sum_{\mu} \frac{1}{2} R^2 \text{Tr} \phi \Gamma_{\mu} \phi \Gamma_{\mu}^{\dagger} - \frac{1}{2} m^2 \phi^2 - \text{Tr} V_0(\phi) \\
V_0(\phi) &= \frac{g_3}{\sqrt{N}} \phi^3 + \frac{g_4}{N} \phi^4 + \dots \quad m^2 = 2DR^2 + m_0^2.
\end{aligned} \tag{2.14}$$

The hopping term is the first term in $S(\phi)$. Adding the effect of world-sheet curvature is easy, we simply replace the hopping term according to

$$\text{Tr}(\Gamma_{\mu} \phi \Gamma_{\mu}^{\dagger} - \phi)^2 \mapsto \text{Tr} A(\Gamma_{\mu} \phi \Gamma_{\mu}^{\dagger} - \phi) A(\Gamma_{\mu} \phi \Gamma_{\mu}^{\dagger} - \phi) \tag{2.15}$$

Our proposal to describe strings in dimensions higher than one is to investigate the properties of (2.14) and (2.15). We have effectively reduced the problem to a one-matrix model. The difficulty lies however in the presence of the twist matrices. We begin our analysis in the following section.

3. PROPERTIES OF THE REDUCED STRING ACTION

Some general features of (2.14) and (2.15) are easy to extract. If the hopping term is very small, we are in the limit where the space-time lattice points are very far apart and the whole surface is mapped into a single point, or various points incoherently. This is the pure gravity regime, γ_{st} is expected to be $-1/2$ and the induced metric plays no role. As the hopping term is increased, it becomes more likely for the surface to occupy more and more space-time lattice sites, the induced metric and singular embeddings in principle begin to play important roles, and we can expect a transition to a phase different from pure gravity at some critical value of the hopping coupling R^2 . We will show presently that in the most naïve approximation, we are driven to a branched polymer phase with $\gamma_{\text{st}} = 1/2$, and at the transition point $\gamma_{\text{st}} = 1/3$. A more careful analysis will show a much broader spectrum of possibilities. Note that the role played by R is similar to the compactification radius in the study of $c = 1$ strings propagating in a circle of radius R [26].

To deal with (2.14) we decompose the matrix ϕ into its eigenvalues and angular variables as usual

$$\phi = U^{-1} \lambda U \quad \lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_N),$$

The angular integration produces an effective action

$$e^{\Gamma_{\text{eff}}(\phi)} = \int dU e^{R^2 \sum_{\mu} \text{Tr} \lambda \Gamma_{\mu}(U) \lambda \Gamma^{\dagger}(U)}$$

$$\Gamma_{\mu}(U) = U \Gamma_{\mu} U^{-1}$$

$$Z = \int d\phi e^{\Gamma_{\text{eff}}[\phi] - V_0(\phi)} \quad (3.1)$$

The simplest representation of Γ_{eff} is given in terms of the leading order expansion in R^2 :

$$Z = \int d\phi e^{N(R^2 \frac{D}{N} (\text{Tr} \phi)^2 - V_0(\phi))}. \quad (3.2)$$

The critical properties of (3.2) are very similar to those of the curvature model in [16]. What the first term in Γ_{eff} is describing is the breaking of the surface. The Feynman graph representation of a term like $(\text{Tr} \phi^n)^2$ is a vertex where two surfaces osculate at one point, and the coordination number of the touching vertices in each of the surfaces is n . If we have instead a term of the form $(\text{Tr} \phi^n)^p$, it represents p -surfaces sharing one point. When there are no products of traces, the Feynman graph expansion for the free energy or the connected correlators involves only the sum over connected orientable surfaces. When products of traces are included we obtain an expansion where we have surfaces of different topologies touching at points determined by the new vertices. The new graphs look like “daisies”. In the original curvature model introduced in [16] the initial partition function is

$$Z = \int d\phi e^{-N(\frac{1}{2} \phi A \phi A + g \text{Tr} \phi^4)} \quad (3.3)$$

The angle integral is again very difficult to carry out in this model, and the simplest

non-trivial approximation is to study the action:

$$S_{\text{eff}} = N \left(\frac{1}{2} \text{Tr} \phi^2 + g \text{Tr} \phi^4 + \frac{g'}{N} (\text{Tr} \phi^2)^2 \right).$$

For real A , the coupling $g' = (\text{Tr} A^2/N - (\text{Tr} A)^2/N^2)$ is always positive. In this case as we will see later one cannot escape the pure gravity phase. To leading order in $1/N$, the graphs have the topology of spheres touching at one point, where any two spheres in the graph can share at most one common point, and no closed loop of spheres can appear. A generic graph looks like a tree of spheres. In the planar approximation it was shown in [16] that we have the following possibilities:

- 1). $g' > 0$, $\gamma_{\text{st}} = -1/2$ and we are in the pure gravity regime.
- 2). $g_c < g' < 0$. Again $\gamma_{\text{st}} = -1/2$ and we continue in the pure gravity case. The branching is still unimportant, and the leading contributions to the free energy come from graphs with few big spheres and few small branchings.
- 3). At $g' = g_c = -9/256$, $\gamma_{\text{st}} = +1/3$ and the number of branchings become competitive with the number of smooth spheres.
- 4). For $g < g_c$ we obtain $\gamma_{\text{st}} = +1/2$ and the dominating term is the one with a product of two traces, thus the notion of a smooth surface disappears and we end up with a branch polymer phase.

The same analysis can be carried out with our naïve approximation (3.2), and we will only note the differences with respect to the analysis in [16] we have just reviewed. The basic differences are:

- 1). Our potential is cubic, $V_0 = \frac{1}{2} \phi^2 + g \phi^3 / \sqrt{N}$, the one-cut solution is asymmetrical, and the new vertex is just given by $\text{Tr}(\phi)^2$. These are purely technical differences and nothing in the analysis changes substantially.
- 2). We are always in the $g' < 0$ region. In the original model in [16] we can reach this region only after analytically continuing the matrix $A \mapsto iA$ in order to obtain the polymer phase. However, in this case the interpretation of the matrix A in terms of local world-sheet curvature loses its meaning.

Our simplified model then contains a pure gravity phase, and intermediate phase with $\gamma_{\text{st}} = 1/3$ and a branched polymer phase with $\gamma_{\text{st}} = 1/2$. We should like to remark that the exact solution to the pure curvature model (3.3) is known only for Penner's potential [27], however in this case the full theory has the symmetry $A \mapsto \text{const.} A^{-1}$ making the interpretation of A in terms of world-sheet curvature rather doubtful.

4. PRELIMINARY ANALYSIS OF THE EFFECTIVE ACTION

We have learned that the basic problem is the evaluation of the angular integral

$$e^{\Gamma_{\text{eff}}} = \int dU e^{R^2 \sum_{\mu} \text{Tr} \lambda U \Gamma_{\mu} U^{-1} \lambda U \Gamma_{\mu}^{\dagger} U^{-1}}. \quad (4.1)$$

The naïve approximation embodied in (3.2) already gave us a polymer phase and a critical point in-between pure gravity and polymers. To understand the behavior of strings in dimensions higher than one, we need to obtain information about the general critical properties of the effective action in (4.1). Since R^2 is a hopping parameter, we can write the hopping term as $L_{\mu} = \phi \Gamma_{\mu} \phi \Gamma_{\mu}^{\dagger}$ and in analogy with standard lattice analysis, we could write an approximation to the effective action in terms of a sum over connected graphs of angular averages over the hopping terms generating the graph:

$$\Gamma_{\text{eff}} = \sum_{C, \text{Connected}} \langle \text{Tr} \prod_{l \in C} L(l) \rangle_U,$$

where the subscript U indicates that the average is taken only over the angular variables. To every connected graph the average over angles associates a set of product of traces. We can interpret these traces as different ways of fracturing and breaking the loop C . They describe how the embedded surface is collapsed into pieces. If we add a local curvature term (the A -matrix), we will obtain more general operators. Since we do not have an explicit way of evaluating (4.1) and

since we are interested in the critical behavior of (2.14) and not in the fine details of lattice dynamics, it is worth exploring the type of universality classes of potentials with arbitrary numbers of products of traces. There are two qualitative classes of potentials, depending on whether they contain a finite or infinite number of traces. So far we have analyzed the case of finite numbers of traces. Defining

$$x_i = \frac{1}{N} \text{Tr} \phi^{2i}, \quad (4.2)$$

we take the effective action to be a general function of an arbitrary, but finite number of x_i variables,

$$\Gamma_{\text{eff}} = V(x_1, x_2, \dots, x_n) = V\left(\frac{1}{N} \text{Tr} \phi^2, \frac{1}{N} \text{Tr} \phi^4, \dots, \frac{1}{N} \text{Tr} \phi^{2n}\right). \quad (4.3)$$

Thus we are faced with the analysis of the critical properties in the planar limit of the action

$$S(\phi) = N^2 V(x_i) + N \text{Tr} V_0(\phi)$$

$$V_0 = \sum_k g_k \text{Tr} \phi^{2k}. \quad (4.4)$$

The world-sheet cosmological constant is related to the coupling $g_2 = e^{-\mu_B}$. We have made the simplifying assumption of restricting our considerations to the case of even potentials. This is just a question of technical simplicity, and nothing changes if we take general potentials. the analysis can be carried out for any of the Kazakov potentials [28]. The solution of the planar approximation of (4.4) uses the Hartree-Fock approximation, whi becomes exact in the planar limit. We first write down the planar equations for (4.4), then, we replace the traces (4.2) by arbitrary variables x_i . This reduces the problem to the original pure gravity case studied in [21], and then we fix the variables x_i self-consistently. The saddle-point equations

and the Hartree-Fock conditions are given by

$$\frac{1}{2} \sum_p 2p \tilde{g}_p \lambda^{2k-1} = \text{P.V.} \int d\mu \frac{\rho(\mu)}{\lambda - \mu}$$

$$\tilde{g}_p = g_p + \frac{\partial V(x)}{\partial x_k} \quad (4.5)$$

$$x_k = \int \rho(\mu) \mu^{2k}. \quad (4.6)$$

$\rho(\mu)$ is the density of eigenvalues. We will denote by $\tilde{V}$ the potential with the shifted couplings (4.5). Following the pure gravity case, we look for one-cut solutions of (4.5). This means that the loop operator

$$F(p) = \int d\mu \frac{\rho(\mu)}{p - \mu}$$

is given by

$$F(p) = \frac{1}{2} \tilde{V}'(p) - M(p) \sqrt{p^2 - R},$$

and the polynomial $M(p)$ is determined by requiring that at large values of $|p|$, $F(p) \sim 1/p + \mathcal{O}(1/p^3)$. Since V depends only on a finite number of traces, the potential $\tilde{V}$ is a polynomial, and there are no ambiguities in the determination of M . Expanding $F(p)$ in powers of $1/p$ we obtain the expectation values of the traces x_i ,

$$F(p) = \sum_{k \geq 0} \frac{x_k}{p^{2k+1}} = \frac{1}{p} + \frac{x_1}{p^3} + \dots \quad (4.7)$$

and the first condition $x_0 = 1$ is the string equation. From the other coefficients we can read off the Hartree-Fock conditions (4.6). The density of eigenvalues is

given by

$$\rho(\lambda) = \frac{1}{2\pi} M(\lambda) \sqrt{R - \lambda^2} \quad \lambda \in [-\sqrt{R}, \sqrt{R}]$$

The dependence of x_k on R is obtained following standard arguments in the planar limit, and we obtain

$$x_p(R) = \sum_{k \geq 1} \frac{k^2}{k+p} \binom{2p}{p} \binom{2k}{k} \left(\frac{R}{4}\right)^{k+p} \tilde{g}_k. \quad (4.8)$$

The string equation is given by (4.8) with $p = 0$. We are interested in the analysis of the critical points of (4.4), the computation of string susceptibilities, and in the computation of correlation functions of macroscopic loop operators. We present only the results, the details will appear elsewhere [29]. If we denote by β the coupling g_2 , the string susceptibility is determined from

$$\chi = \frac{dx_2}{d\beta} = \frac{\partial x_2}{\partial \beta} + \frac{dR}{d\beta} \frac{\partial x_2}{\partial R}. \quad (4.9)$$

This is because the coupling β multiplies the operator x_2 in the action. Therefore $\partial \ln Z / \partial \beta$ is proportional to the expectation value of x_2 , and the second derivative of the logarithm of the partition function is the string susceptibility. The dependence of R on β is read off from the string equation $x_0 = 1$. In particular,

$$0 = \frac{dx_0}{d\beta} = \frac{\partial x_0}{\partial \beta} + \frac{dR}{d\beta} \frac{\partial x_0}{\partial R}$$

allows us to read the equation determining the critical points,

$$\frac{d\beta}{dR} = -\frac{\partial_R x_0}{\partial_\beta x_0}, \quad (4.10)$$

Since the denominator in (4.10) is proportional to R^2 , the critical points are determined by the zeroes of the numerator. We can similarly write χ according to

$$\chi = \partial_\beta x_2 + \partial_\beta x_0 \frac{\partial_R x_2}{\partial_R x_0} \quad (4.11)$$

The singularities in the behavior of χ come from the ratio of derivatives in (4.11). For the standard Kazakov critical points [28] (4.11) does not blow up at the critical

point R_c , and $\chi - \chi_c \sim R - R_c$, however at the m -th critical point, $\beta - \beta_c \sim (R - R_c)^m$, and this yields $\gamma_{\text{st}} = -1/m$. As we will show presently, the critical points we describe have $\gamma_{\text{st}} > 0$ and χ will be singular at the critical point. All the phase transitions are second order.

In analyzing (4.5), (4.6) it is useful to distinguish between two types of derivatives. When we differentiate with respect to R , we can consider first the explicit dependence of the equations without taking into account the Hartree-Fock condition. If we ignore the implicit dependence on R of $\tilde{g}_k$ we call this derivative D_R , and everything is as in the pure gravity case. It is not difficult to show that

$$D_R x_p(R) = \binom{2p}{p} \left(\frac{R}{4}\right)^p D_R x_0(R). \quad (4.12)$$

Including the R dependence coming from (4.6) we have an extra contribution. The derivatives with respect to R including the Hartree-Fock induced dependence on R will be denoted by ∂_R . Thus,

$$\begin{aligned} \partial_R X_p &= D_R x_p + \sum_{k \geq 1} \frac{k^2}{k+p} \binom{2k}{k} \binom{2p}{p} \left(\frac{R}{4}\right)^{k+p} \partial_R \tilde{g} \\ &= D_R x_p + \sum_{k \geq 1} \frac{k^2}{k+p} \binom{2p}{p} \left(\frac{R}{4}\right)^p \binom{2k}{k} \left(\frac{R}{4}\right)^k \sum_q \partial_{pq} V \partial_R x_q. \end{aligned} \quad (4.13)$$

Defining the matrix

$$U''_{pq} = \sum_{k \geq 1} \frac{k^2}{k+p} \binom{2p}{p} \left(\frac{R}{4}\right)^p \binom{2k}{k} \left(\frac{R}{4}\right)^k \partial_{pq} V, \quad (4.14)$$

we can compute $\partial_R x_p$ in terms of $D_R x_0$:

$$\partial_R x_p = [(1 - U'')^{-1} A]_p D_R x_0, \quad (4.15)$$

where A is a vector whose p -th component equals $\binom{2p}{p} \left(\frac{R}{4}\right)^p$. When $p = 0$, we

obtain the criticality condition implied by (4.10):

$$\partial_R x_0 = [(1 - U'')^{-1} A]_0 D_R x_0 = 0. \quad (4.16)$$

Notice that the criticality condition splits into two terms. The first one is related to the function including the couplings with multiple traces,

$$[(1 - U'')^{-1} A]_0 = 0, \quad (4.17)$$

and the second is equivalent to the criticality condition for the Kazakov critical points:

$$D_R x_0 = 0. \quad (4.18)$$

The term in (4.11) which may lead to singularities in χ and to positive γ_{st} is given by

$$\frac{\partial_R x_2}{\partial_R x_0} = \frac{[(1 - U'')^{-1} A]_2}{[(1 - U'')^{-1} A]_0}. \quad (4.19)$$

In (4.16) we can have three possibilities: 1). The polymer couplings in U become critical but the gravity part does not. The zeroes of $d\beta/dR$ come only from the polymer contribution. 2). The gravity contribution is the only one generating zeroes of $d\beta/dR$. 3). Both terms becomes critical. In the first case we have a theory of polymers, and little is remembered of the coupling to gravity. In the second case the polymer degrees of freedom are completely frozen and we reproduce the string susceptibility and exponents of the Kazakov critical points. The third and more interesting case is when both the “polymer” matter and gravity become critical simultaneously. This is the case where we can find novel behavior. Notice also that unless we tune the numerator in (4.19) we will generically obtain $\gamma_{\text{st}} > 0$ in cases 1) and 3). If R_c, β_c are the critical values of R and β , and if near the

critical point

$$[(1 - U'')^{-1}A]_0 \sim (R - R_c)^n \quad D_R x_0 \sim (R - R_c)^m, \quad (4.20)$$

then

$$\beta - \beta_c \sim (R - R_c)^{n+m+1}. \quad (4.21)$$

If we do not tune the numerator of (4.19) to partially cancel the zeroes in the denominator, we obtain

$$\chi \sim \frac{1}{(R - R_c)^n}.$$

The previous two equations give a string susceptibility for the (n, m) -critical point:

$$\gamma_{\text{st}} = \frac{n}{n + m + 1}. \quad (4.22)$$

By tuning the numerator in (4.19) we can change the numerator in (4.22) to any positive integer smaller than n

$$\gamma_{\text{st}} = \frac{p}{n + m + 1}.$$

We will show below that only in the case $n = 1$ the model maintains some resemblance with the properties one would expect of a non-critical string. We can define operators creating macroscopic loops only in that case. For $n > 1$ the polymerization of the surface is so strong that there is no room left to open macroscopic loops. The simplest case to study is the one where the function V depends on a single trace $V = V(x_l)$. In this case the matrix U'' becomes:

$$U''_{pl} = \frac{l^2}{l + p} \binom{2p}{p} \left(\frac{R}{4}\right)^p \binom{2l}{l} \left(\frac{R}{4}\right)^l V'' \quad p = 1, 2, \dots$$

The matrix U'' contains a single non-vanishing column in position l . The inverse

of $1 - U''$ is:

$$(1 - U'')^{-1} = 1 + \frac{U''}{1 - \frac{l}{2} \binom{2l}{l} \left(\frac{R}{4}\right)^l V''},$$

therefore,

$$\begin{aligned} [(1 - U'')^{-1} A]_p &= \\ \binom{2p}{p} \left(\frac{R}{4}\right)^p &+ \frac{1}{1 - \frac{l}{2} \binom{2l}{l} \left(\frac{R}{4}\right)^l V''} \frac{l^2}{p+l} \binom{2l}{l} \left(\frac{R}{4}\right)^l \binom{2p}{p} \left(\frac{R}{4}\right)^p V'' \\ [(1 - U'')^{-1} A]_0 &= \frac{1 + \frac{l}{2} \binom{2l}{l} \left(\frac{R}{4}\right)^l V''}{1 - \frac{l}{2} \binom{2l}{l} \left(\frac{R}{4}\right)^l V''}. \end{aligned} \quad (4.23)$$

From these formulae we can immediately derive the string susceptibility χ . Without loss of generality we can restrict our considerations to the case when $l = 2$. In this case many of the equations simplify and all the main results remain the same. The one-point functions x_0, x_2 are given by

$$\begin{aligned} 1 = x_0(R) &= \sum_{k \geq 1} k \binom{2k}{k} \left(\frac{R}{4}\right)^k g_k + 12V'(x_2) \left(\frac{R}{4}\right)^2 \\ x_2 &= \sum_{k \geq 1} \frac{6k^2}{k+2} \binom{2k}{k} \left(\frac{R}{4}\right)^{k+2} g_k + 36V'(x_2) \left(\frac{R}{4}\right)^4. \end{aligned} \quad (4.24)$$

Restricting the general formulae derived before to this case we obtain:

$$x_2 = 3 \left(\frac{R}{4}\right)^2 + 3 \sum_{k \geq 1} \frac{k(k-2)}{k+2} \binom{2k}{k} \left(\frac{R}{4}\right)^{k+2} g_k$$

$$\partial_R x_2 = \frac{6 \left(\frac{R}{4}\right)^2}{1 - 36 \left(\frac{R}{4}\right)^4 V''} D_R x_0$$

$$RD_R x_0 = 2 + \sum_{k \geq 1} k(k-2) \binom{2k}{k} \left(\frac{R}{4}\right)^k g_k. \quad (4.25)$$

The critical points come from the vanishing of the last equation and from

$$1 + 36 \left(\frac{R}{4}\right)^4 V'' = 0.$$

An argument similar to the one presented at the end of [16] shows that the critical point for the transition between the polymer and the pure gravity phases is generically second order, although by tuning of the parameters it could be made of higher order. Hence we can define a continuum limit for the critical points (4.22).

To explore the new critical points in some detail, we compute the correlation functions of macroscopic loop operators. The simplest loops to compute are the ones analogous to those appearing in pure gravity (see for instance [30]). They are obtained by taking the limit $k \rightarrow \infty$ of $Tr\phi^{2k}$. In terms of the original reduced model these are loop on the surface whose position is averaged over the target space. One could also define loops with definite positions in the target space. The expectation values of the latter would certainly give crucial information on the reduced formulation of string theory, but we have not yet calculated them. For the simpler, pure gravity loops of length l we take the definition

$$w(l) = \lim_{k \rightarrow \infty} \sqrt{\frac{\pi}{l}} \langle Tr\phi^{2k} \rangle = N \lim_{k \rightarrow \infty} \sqrt{\frac{\pi}{l}} x_k. \quad (4.26)$$

As a matter of convenience we choose the critical values for β and R as $\beta_c = -1$, $R_c = 1$. In the Kazakov critical points, $\gamma_{st} = -1/m$ and the scaling limit is taken to be $Na^{2-\gamma_{st}} = \kappa^{-1}$. It is a simple but non-trivial fact that the correct way to define a loop of length l in this case is to scale k according to $ka^{-2\gamma_{st}} = l$ as $k \rightarrow \infty$ (see for example [30] and references therein). In our case we will restrict for simplicity our computations to the case of potentials $V = V(x_2)$. The relevant

formulae to be used are (4.23-25). We have in general that

$$\frac{dx_k}{d\beta} = \partial_\beta x_k - \partial_\beta x_0 \frac{\partial_R x_k}{\partial_R x_0}, \quad (4.27)$$

and from

$$\partial_\beta x_k = \frac{24}{k+2} \binom{2k}{k} \left(\frac{R}{4}\right)^{k+2} (1 + V'' \partial_\beta x_2)$$

we obtain

$$\partial_\beta x_k = \frac{2}{k+2} \binom{2k}{k} \left(\frac{R}{4}\right)^k \partial_\beta x_0. \quad (4.28)$$

Simple manipulations yield:

$$\frac{\partial_R x_k}{\partial_R x_0} = \binom{2k}{k} \left(\frac{R}{4}\right)^k \frac{1 + 36 \frac{2-k}{2+k} \left(\frac{R}{4}\right)^4 V''}{1 + 36 \left(\frac{R}{4}\right)^4 V''}. \quad (4.29)$$

and after some algebra we obtain:

$$\frac{dx_k}{d\beta} = \frac{12k}{k+2} \binom{2k}{k} \left(\frac{R}{4}\right)^{k+2} \frac{1}{1 + 36 \left(\frac{R}{4}\right)^4 V''} \quad (4.30)$$

Up to irrelevant numerical factors this answer coincides with the one-loop expectation value in the Kazakov critical points when $V'' = 0$. To go to the continuum limit we have to choose the scaling variables. Recall that in the double scaling limit [11] the free energy takes the form

$$F = \sum_{g \geq 0} N^{2-2g} (\mu - \mu_c)^{(2-\gamma_{\text{st}})(1-g)} F_g$$

where $\mu = -\log \beta$, and g is the genus of the triangulations summed over. The renormalized cosmological constant is introduced according to $\mu - \mu_c = a^2 t$ where a is the microscopic lattice spacing. Then the effective string coupling constant is defined by $N a^{2-\gamma_{\text{st}}} = \kappa^{-1}$. Hence the scaling variable associated with the string

susceptibility is $\chi = a^{-2\gamma_{\text{st}}}u$. At the (n, m) -critical point χ behaves as $\chi \sim (1 - R)^{-n}$. Thus the scaling variables are

$$\chi \sim \frac{1}{(1 - R)^n} = a^{-2\gamma_{\text{st}}}u \quad 1 - R = a^{2\gamma_{\text{st}}/n}u^{-1/n}$$

$$\beta - \beta_c = \beta + 1 = a^2t. \quad (4.31)$$

For an (n, m) -critical point we have

$$1 - R = a^{\frac{2}{n+m+1}}u^{\frac{-1}{n}} \quad \chi = ua^{\frac{2n}{n+m+1}} \quad Na^{2-\frac{n}{n+m+1}} = \kappa^{-1}. \quad (4.32)$$

Let us start with the critical points $(1, m)$. From (4.30), and (4.32) we easily obtain as $k \rightarrow \infty$

$$\frac{d}{dt}\langle Tr\phi^{2k} \rangle = -\frac{3}{16} \frac{Na^{2-\frac{2}{m+2}}}{\sqrt{\pi k}} u e^{-ka^{\frac{2}{m+2}}u^{-1}}, \quad (4.33)$$

thus, choosing $ka^{2/m+2} = l$ fixed as $k \rightarrow \infty$ and $a \rightarrow 0$ together with (4.32) we obtain a finite result

$$\frac{d}{dt}w(l) = -\frac{3u}{16\kappa l} e^{-l/u}. \quad (4.34)$$

If we take instead $n > 1$, it is easy to show that there is no possible definition of the macroscopic length l giving a finite result for (4.33). When $n > 1$ we have

$$1 + 36 \left(\frac{R}{4} \right)^4 V'' \sim (1 - R)^n.$$

Repeating previous arguments we arrive at

$$\frac{d}{dt}\langle Tr\phi^{2k} \rangle = -\frac{3}{4} Na^{2-\frac{2n}{m+n+1}} \frac{1}{\sqrt{\pi k}} e^{-ka^{\frac{2}{m+n+1}}u^{-1/n}},$$

and there is no way of choosing the scaling behavior of k giving a finite value without extraneous renormalization of the loop operator. Hence in the phases

with $n > 1$ the polymer couplings dominate completely the critical behavior in spite of the fact that gravity becomes also critical in such a way that there is not enough room to open macroscopic loops. In the case of $n = 1$ we can also calculate two- and multiloop correlation functions. The result obtained for the two-loop correlators is rather suggestive,

$$w(l_1, l_2) = \frac{e^{-(l_1+l_2)/u}}{l_1 + l_2} + \frac{u}{l_1 l_2} e^{-(l_1+l_2)/u} \quad (4.35)$$

and similar results for multiloop correlators. The lesson we can draw from (4.35) is that in the simple case of $n = 1$ the effect of the polymer couplings is to contribute an extra state (the last term in (4.35)) which resembles very much the contribution one would expect of a tachyon. This term represents the breaking of the cylinder interpolating between the two loop of lengths l_1, l_2 into two disks osculating at one point. We can represent the last term in (4.35) as

$$\langle w(l_1)P \rangle \frac{1}{\langle PP \rangle} \langle Pw(l_1) \rangle$$

where P is the puncture operator. This pattern repeats in the higher loop planar correlators, and it is reminiscent of the residue of the tachyon pole in the Belavin-Knizhnik theorem [31]. This interpretation, though tempting has to be taken with several grains of salt because in the types of effective actions we are studying, it is hard to see any dependence on dimensionality. Nevertheless we find this result as encouraging evidence that our approximation captures some of the expected properties of non-critical strings for $D > 1$.

The form of the two-loop operator (4.35) does not depend on the form of the potential $V(x_1, \dots, x_n)$ as long as we consider $(1, m)$ -critical points. Hence only the quadratic part of V determines the critical properties of macroscopic loops. Whether these conclusions will still hold when we consider potentials V depending on an infinite number of traces is currently under investigation [29].

To conclude this long section we mention that we can study at least in part the spectrum of scaling operators by perturbing the criticality conditions. The string equation is then modified according to

$$1 + \beta = (1 - R)^{n+m+1} + \sum t_k^B (1 - R)^{k+1}$$

The subleading contributions depend on both the polymer and Kazakov perturbations of the critical conditions (4.16). The bare couplings of the scaling operators are the t_k^B parameters. Using (4.32) the string equation becomes

$$t = u^{-\frac{n+m+1}{n}} + \sum_k t_k^R u^{-\frac{k+1}{n}}, \quad (4.36)$$

and the t_k are the renormalized couplings. In the original model these couplings would hardly exhaust the scaling operators, although they may represent a significant subset associated to the simplest loop operators. If we call σ_k the scaling operator associated to the coupling t_k , its planar correlators at the (n, m) -critical point follow from (4.36)

$$\frac{du}{dt_k} = -u^{-\frac{k+1}{n}} \frac{du}{dt}$$

and

$$\langle \sigma_k PP \rangle = \frac{n}{n+m+1} t^{\frac{k-m-2n}{n+m+1}}.$$

This concludes our brief study of the effective actions depending on an arbitrary, but finite number of couplings.

5. CONCLUSIONS AND OUTLOOK

We have presented a very preliminary analysis of a new way of formulating strings theories in dimensions higher than one. Our approach is a direct application of the reduced formulation of large- N field theories [1,2]. We believe that it is worth while exploring further the properties of these reduced models to gain insight into the behavior of strings beyond $D = 1$.

We have presented the general analysis of effective actions in the planar limit containing arbitrary, but finite numbers of traces (4.4). We were able to show that beyond some critical coupling, γ_{st} becomes positive and takes values of the form $\gamma_{\text{st}} = n/n + m + 1$ for any positive integers n, m . We should continue exploring the properties of these critical points and their scaling operators. Obviously the more challenging case of studying arbitrary functions V depending on an infinite number of traces is crucial before we can draw any conclusions concerning the properties of our model. This together with the study of general properties of (4.1) should shed further light into the properties of non-critical strings in interesting dimensions $D = 2, 3, 4$. It is very important to obtain in what way the effective actions studied in the previous sections depend on dimensionality.

To understand better the transition between $\gamma_{\text{st}} < 0$ and $\gamma_{\text{st}} > 0$ in the models analyzed, we calculated the correlation functions of several macroscopic loops of lengths $l_1, l_2, \dots$. We found that only for critical points of the form $(1, m)$ is it possible to open macroscopic loop on the surface. For $n > 1$ the polymer couplings dominate the continuum limit and there is no room to open such loops. Furthermore in the case of $n = 1$ we encountered a new state which represents the breaking of the surface into two pieces touching only at one point, qualitatively in agreement with what one would expect from the presence of a tachyon beyond $D = 1$. Whether this conclusion will remain after one includes more realistic effective actions remains to be seen. However this “stringy” aspect of our effective action seems encouraging. We can also compute in the reduced models the explicit form of ordinary vertex operators which describe embedded surfaces passing

through specified target space points. Correlation functions of vertex operators should allow us to better assess the quantum geometry of the phases for $\gamma_{\text{st}} > 0$. Furthermore, the study of vertex operators should also help in the interpretation of the operators $\prod_n \text{Tr} \phi^n$ in terms of processes tearing up the embedded surface and the effect of tachyons. Finally it may also be useful to analyze strings in $D > 1$ numerically using the reduced actions.

Many of these issues are currently under investigation. Details concerning the topics covered in this lecture together with further results will appear elsewhere [29].

ACKNOWLEDGEMENTS. One of us (L. A.-G.) would like to thank the organizers of the Trieste Spring Workshop 1992 for the opportunity to present these results in such a stimulating environment.

REFERENCES

1. T. Eguchi and H. Kawai, Phys. Rev. Lett. **48** (1982) 1063.
2. A. Gonzalez-Arroyo and M. Okawa, Phys. Rev. **D 27** (1983) 2397, Phys. Lett. **B 133**(1983) 415, Nucl. Phys. **B 247** (1984) 104.
3. For a review with references to the literature see S.R. Das, Rev. Mod. Phys. **59** (1987) 235.
4. J. Ambjörn, B. Durhuus and J. Fröhlich, Nucl. Phys. **B 259** (1985) 433.
5. J. Ambjörn, B. Durhuus and J. Fröhlich and P. Orland, Nucl. Phys. **B 270** (1986) 457.
6. F. David, Nucl. Phys. **B 257** (1985) 53, Nucl. Phys. **B 257** (1985) 543.
7. V.A. Kazakov, Phys. Lett. **150 B** (1985) 282.
8. A.M. Polyakov, Mod. Phys. Lett. **A2** (1987) 893; V. G. Knizhnik, A.M. Polyakov and A.B. Zamolodchikov, Mod. Phys. Lett. **A 3** (1988) 819.
9. F. David, Mod. Phys. Lett. **A 3** (1988) 1651; J. Distler and H. Kawai, Nucl. Phys. **B 321** (1988) 509.
10. V.A. Kazakov, Phys. Lett. **119 B** (1986) 140, D. Boulatov and V. A. Kazakov, Phys. Lett. **186 B** (1987) 379; I.K. Kostov, and M.L. Mehta, Phys. Lett. **189 B** (1987) 118; B. Duplantier and I. Kostov, Phys. Rev. Lett. **61** (1988) 1433; etc.
11. M.R. Douglas, S. Shenker, Nucl. Phys. **B 335** (1990) 635; E. Brézin and V.A. Kazakov, Phys. Lett. **236 B** (1990) 144; D. Gross and A.A. Migdal, Phys. Rev. Lett. **64** (1990) 127, Nucl. Phys. **B 340** (1990) 333.
12. See J.-L. Gervais, Comm. Math. Phys. **138** (1991) 301; Phys. Lett. **243 B** (1990) 85, and references therein.
13. E. Brezin and D. Gross, Phys. Lett. **97 B** (1980) 120.
14. D. Gross and Newman, Phys. Lett. **266 B** (1991) 291; PUPT-1282, Dec. 91.

15. E. Witten, Surveys in Diff. Geom. **1** (1991) 243, *On the Kontsevich Model and Other Models of Two Dimensional Gravity*. IAS preprint HEP-91/24. M. Kontsevich, Funct. anal. i ego pril. **25** (1991) 50; *Intersection Theory on the Moduli Space of Curves and the Matrix Airy Function*. M. Plank Institute for Mathematics Preprint, October 91.
16. S.R. Das, A. Dhar, A.M. Sengupta and S.R. Wadia, Mod. Phys. Lett. **A5** (1990) 1041.
17. A.M.Polyakov, Phys. Lett. **103 B** (1981) 207, 211.
18. D. Boulatov, V.A. Kazakov, I. Kostov, A.A. Migdal, Nucl. Phys. **B 275** (1986) 641.
19. D. Boulatov, V.A. Kazakov, Phys. Lett. **214 B** (1988) 581; J. Ambjörn, Acta Phys. Pol. **B21** (1990) 101, and references therein.
20. E. Witten, *Recent Developments in Gauge Theories*, Proceedings of the 1979 Cargèse Summer Institute. G. 'tHooft et al. eds. Plenum Publ. New York.
21. E. Brézin C. Itzykson, G. Parisi and J.B. Zuber, Comm. Math. Phys. **59** (1979) 35.
22. Yu. M.Makeenko, and A.A. Migdal, Phys. Lett. **88 B** (1979) 135; S. Wadia, Phys. Rev. **D 24** (1981) 970. For more details and references see A.A. Migdal, Phys. Rep. **102** (1983) 200.
23. G. Bhanot, U. Heller, H. Neuberger, Phys. Lett. **113 B** (1982) 47.
24. G. 'tHooft, Nucl. Phys. **B 153** (1979) 141, Acta Phys. Austriaca, Suppl. **22** (1980) 531.
25. T. Eguchi R. Nakayama, Phys. Lett. **122 B** (1983) 59.
26. D. Gross and I. Klebanov, Nucl. Phys. **B 344** (1990) 475.
27. Yu. Makeenko and L. Chekhov, *The Multicritical Kontsevich-Penner Model*. NBI-HE-92-03, January 1992.
28. V.A. Kazakov, Mod. Phys. Lett. **A4** (1989) 2125.

- 29. L. Alvarez-Gaumé, J.L.F. Barbón and C. Crnkovic, in preparation.
- 30. G. Moore, N. Seiberg and M. Staudacher, Nucl. Phys. **B 362** (1991) 665.
- 31. A.A. Belavin and V. Knizhnik, Phys. Lett. **168 B**(1986) 201; J.B. Bost and T. Jolicoeur, Phys. Lett. **174 B**(1986) 273; R. Catenacci, M. Cornalba, M. Martellini and C. Reina, Phys. Lett. **172 B**(1986) 328.